

PC1134 Lecture 17

Topic


Line integrals

Application example

Work done by a force.

$$dW = \vec{F} \cdot d\vec{r}$$

$$W = \int_A^B \vec{F} \cdot d\vec{r}$$




Scope

- Review of integration
- Line integral
- Calculation of line integral

Integration

The **definite integral** of $f(x)$ between the **lower limit** $x = a$ and the **upper limit** $x = b$. $f(x)$ is called the **integrand**.

$$I = \int_a^b f(x) dx$$




I is the **area** under the curve.

Integration from First Principles

Divide the area into slices of width dx (not required to be uniform). The area of each slice is

$$dA \approx f(x)dx$$



Total area

$$A \approx \sum f(x)\Delta x$$

The error is reduced when Δx is small. In the limit of $\Delta x \rightarrow 0$,

$$A = \lim_{\Delta x \rightarrow 0} \sum f(x)dx = \int_a^b f(x)dx$$

Properties of Integration

$$\int_a^b f(x)dx = - \int_b^a f(x)dx$$

$$\int_a^c f(x)dx = \int_a^b f(x)dx + \int_b^c f(x)dx$$

$$\int_a^b [f(x) + g(x)]dx = \int_a^b f(x)dx + \int_a^b g(x)dx$$

$$\frac{d}{dx} \left[\int_a^x f(t)dt \right] = f(x)$$

Integration of Simple Functions

$$\int a dx = ax$$

$$\int x^n dx = \frac{1}{n+1} x^{n+1}$$

$$\int e^{ax} dx = \frac{e^{ax}}{a}$$

$$\int \frac{dx}{x} = \ln x$$

$$\int \cos ax dx = \frac{\sin ax}{a}$$

$$\int \sin ax dx = -\frac{\cos ax}{a}$$

$$\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right)$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \left(\frac{x}{a} \right)$$

.....

Integration by Substitution

Example 1

$$I = \int \frac{dx}{\sqrt{1-x^2}}$$

Let

$$x = \sin u$$

then

$$dx = \cos u \, du$$

$$I = \int \frac{\cos u \, du}{\sqrt{1 - \sin^2 u}} = \int \frac{\cos u \, du}{\sqrt{\cos^2 u}} = \int du = u$$

$$I = \sin^{-1} x$$

Example 2

$$I = \int \frac{dx}{x^2 + 4x + 7} = \int \frac{dx}{(x+2)^2 + 3}$$

Let $y = x + 2$, then $dy = dx$ and

$$I = \int \frac{dy}{y^2 + 3} = \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{y}{\sqrt{3}} \right) = \frac{1}{\sqrt{3}} \tan^{-1} \left(\frac{x+2}{\sqrt{3}} \right)$$

Integration by Parts

If u and v are functions of x

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + \frac{du}{dx}v$$

Integrate both sides

$$uv = \int u \frac{dv}{dx} dx + \int \frac{du}{dx} v dx$$


This can be rewritten as

$$\int u \frac{dv}{dx} dx = uv - \int \frac{du}{dx} v dx$$

Example

$$\begin{aligned} I = \int x \sin x dx &= \int x \frac{d(-\cos x)}{dx} dx = x(-\cos x) \\ &\quad - \int (1)(-\cos x) dx = -x \cos x + \sin x \end{aligned}$$

Work Done by a Varying Force



$$dW = \vec{F} \cdot d\vec{r}$$

$$W = \int \vec{F} \cdot d\vec{r}$$

- \vec{F} is a function of position

$$\vec{F}(\vec{r}) \text{ or } \vec{F}(x, y, z)$$

- Object is restricted to move along the curved path.

Line Integral

$$W = \int \vec{F} \cdot d\vec{r}$$

2D

A curve is described by one equation

$$\phi(x, y) = 0 \quad \Longrightarrow \quad y = \psi(x)$$

$$\int \vec{F}(x, y) \cdot d\vec{r} \quad \Longrightarrow \quad \int f(x) dx$$

or

$$\int \vec{F}(x, y) \cdot d\vec{r} \quad \Longrightarrow \quad \int f(s) ds$$

3D

Two equations relating x , y and z are required to describe a curve

\Longrightarrow Only one independent variable

\Longrightarrow One dimensional integral

Example


$$\vec{F} = xy\hat{x} - y^2\hat{y}$$

Calcualte

$$W = \int \vec{F} \cdot d\vec{r}$$


along

1. Path 1 (black) $y = x/2$
2. Path 2 (red) $y = x^2/4$
3. Path 3 (green) $x = 0$ and $y = 1$
4. Path 4 (blue) $x = 2t^3$, $y = t^2$.



Example

Path 1 (black)



$$y = \frac{1}{2}x, \quad dy = \frac{1}{2}dx$$

$$d\vec{r} = dx\hat{x} + dy\hat{y} = dx\hat{x} + \frac{1}{2}dx\hat{y} = \left(\hat{x} + \frac{1}{2}\hat{y}\right) dx$$


$$\vec{F} \cdot d\vec{r} = F_x dx + F_y dy = xy dx - y^2 \frac{1}{2} dx = \left(xy - \frac{1}{2}y^2\right) dx$$

$$= \left(\frac{1}{2}x^2 - \frac{1}{8}x^2\right) dx = \frac{3}{8}x^2 dx$$

$$W_1 = \int \vec{F} \cdot d\vec{r} = \int_0^2 \frac{3}{8}x^2 dx = \frac{x^3}{8} \Big|_0^2 = 1$$

Example

Path 2 (red)



$$y = \frac{1}{4}x^2, \quad dy = \frac{1}{2}x dx$$

$$d\vec{r} = dx\hat{x} + dy\hat{y} = dx\hat{x} + \frac{1}{2}x dx\hat{y} = \left(\hat{x} + \frac{1}{2}x\hat{y}\right) dx$$


$$\vec{F} \cdot d\vec{r} = F_x dx + F_y dy$$

$$= \left(xy - \frac{1}{2}xy^2\right) dx = \left(\frac{1}{4}x^3 - \frac{1}{32}x^5\right) dx$$

$$W_2 = \int \vec{F} \cdot d\vec{r} = \int_0^2 \left(\frac{1}{4}x^3 - \frac{1}{32}x^5\right) dx = \frac{2}{3}$$

Example

Path 3 (green)



Along the vertical segment, $x = 0$ and $dx = 0$

$$\vec{F} \cdot d\vec{r} = F_x dx + F_y dy = -y^2 dy$$

$$W_{3a} = \int \vec{F} \cdot d\vec{r} = \int_0^1 (-y^2) dy = -\frac{1}{3}$$

Along the horizontal segment, $y = 1$ and $dy = 0$

$$\vec{F} \cdot d\vec{r} = F_x dx + F_y dy = x dx$$


$$W_{3b} = \int \vec{F} \cdot d\vec{r} = \int_0^2 x dx = 2$$

$$W_3 = W_{3a} + W_{3b} = -\frac{1}{3} + 2 = \frac{5}{3}$$

Example

Path 4 (blue):

$$x = 2t^3, \quad y = t^2$$



$$\vec{F} = xy\hat{x} - y^2\hat{y} = 2t^5\hat{x} - t^4\hat{y}$$

$$d\vec{r} = dx\hat{x} + dy\hat{y} = 6t^2dt\hat{x} + 2tdt\hat{y}$$

$$\vec{F} \cdot d\vec{r} = F_x dx + F_y dy = 12t^7 dt - 2t^5 dt$$

$$W_4 = \int \vec{F} \cdot d\vec{r} = 12 \int_0^1 t^7 dt - 2 \int t^5 dt = \frac{7}{6}$$

Summary

- $\vec{F}(x, y, z)$ is a function of one variable because x , y and z are related.
- This variable can be one of x , y and z or can be something else.
- The integration over $\vec{F} \cdot d\vec{r}$ is an one-dimensional integral over this variable.
- The line integral between two points can be path dependent. ($W_1 = 1$, $W_2 = 2/3$, $W_3 = 5/3$, $W_4 = 7/6$).