

# Lecture 27

## Topic

Matrix operations

## Relevance

Linear operators and vectors can be represented by matrices. A knowledge of matrix analysis is required.

## Objective

To understand and able to perform matrix addition, subtraction, multiplication, differentiation, integration; to be able to calculate the transpose, comple-conjugate, Hermitian conjugate matrices of a given matrix.

# Addition & Subtraction of Matrices

The operation of **addition** (or **subtraction**) for two  $n \times m$  matrices is defined as

$$C = A \pm B$$

where

$$c_{ij} = a_{ij} \pm b_{ij}$$

Example:

$$\begin{pmatrix} 1 & -1 & 2 \\ 3 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 2 & 2 & 2 \\ 1 & 0 & -1 \end{pmatrix} = \begin{pmatrix} 3 & 1 & 4 \\ 4 & 0 & 0 \end{pmatrix}$$

The **commutative** and **associative laws** are also valid for addition of matrices of the same order:

$$A + B = B + A$$

$$(A + B) + C = A + (B + C)$$

## Equal Matrices & Null Matrix

Two matrices,  $A$  and  $B$ , of the same order are said to be **equal** if and only if  $a_{ij} = b_{ij}$  for all  $i$  and  $j$ . For example,

$$A = \begin{pmatrix} 2 & 0 & -1 & 2 \\ 6 & 5 & 3 & 7 \\ 2 & -1 & 0 & 4 \end{pmatrix}$$

and

$$B = \begin{pmatrix} 2 & 0 & -1 & 2 \\ 6 & 5 & 3 & 7 \\ 2 & -1 & 0 & 4 \end{pmatrix}$$

If  $a_{ij} = 0$  for all  $i$  and  $j$ , then  $A$  is called a **null matrix**. For example

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

## Multiplication of a Matrix by a Scalar

Each element of the matrix is multiplied by the scalar. The elements of  $kA$  are therefore  $ka_{ij}$  for all  $i$  and  $j$ . For example

$$2 \begin{pmatrix} 3 & 1 & 2 \\ 2 & -3 & -1 \\ 1 & 2 & 1 \end{pmatrix} = \begin{pmatrix} 6 & 2 & 4 \\ 4 & -6 & -2 \\ 2 & 4 & 2 \end{pmatrix}$$

## Matrix Product

Matrix product is defined for conformable matrices only. This means that the number of columns of  $A$  must equal the number of rows of  $B$  for  $C = AB$ . Consider

$$\begin{aligned} A &= \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \text{ and } B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \\ AB &= \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix} \\ &= \begin{pmatrix} a_{11}b_{11} + a_{12}b_{21} & a_{11}b_{12} + a_{12}b_{22} \\ a_{21}b_{11} + a_{22}b_{21} & a_{21}b_{12} + a_{22}b_{22} \end{pmatrix}. \end{aligned}$$

In general, the matrix product  $C = AB$  is obtained by use of the following definition:

$$c_{ij} = \sum_{k=1}^l a_{ik}b_{kj}$$

where the orders of  $A$ ,  $B$ , and  $C$  are  $n \times l$ ,  $l \times m$ , and  $n \times m$ , respectively.

## Examples of Matrix Product

$$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix}$$

Consider the following system of equations

$$\begin{array}{rcrcrcrl} 3x & + & y & + & 2z & = & 3 \\ 2x & - & 3y & - & z & = & -3 \\ x & + & 2y & + & z & = & 4 \end{array}$$

In matrix form, it can be written as

$$\begin{pmatrix} 3 & 1 & 2 \\ 2 & -3 & -1 \\ 1 & 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 3 \\ -3 \\ 4 \end{pmatrix}$$

# Matrix Multiplication

Consider

$$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix}$$

$$\begin{pmatrix} 0 & 0 \\ 2 & 3 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 3 \end{pmatrix}$$

It can be shown by use of the definition of the matrix product, the commutative law of multiplication is not, in general, valid for the matrix product,

$$AB \neq BA.$$

However, the associative law of multiplication is valid for the matrix product,

$$A(BC) = (AB)C$$

.

## Transpose Matrix: $A^T$

The **transpose** of an arbitrary matrix  $A$  is written as  $A^T$  and is obtained by interchanging corresponding rows and columns of  $A$ , that is, if the element of  $i$ th row and  $j$ th column of  $A$  is  $a_{ij}$ , then the element at  $i$ th row and  $j$ th column of  $A^T$  is  $a_{ji}$ .

For example

$$A = \begin{pmatrix} 1 & -1 & 2 \\ 3 & 0 & 1 \end{pmatrix}$$

$$A^T = \begin{pmatrix} 1 & 3 \\ -1 & 0 \\ 2 & 1 \end{pmatrix}$$

It can be shown that

$$(AB)^T = B^T A^T$$

$$(ABC \cdots G)^T = G^T \cdots C^T B^T A^T$$



## Complex-Conjugate Matrix: $A^*$

The **complex conjugate** of an arbitrary matrix  $A$  is formed by taking the complex conjugate of each element. Hence we have

$$(A^*)_{ij} = a_{ij}^* \quad \text{for all } i \text{ and } j.$$

For example

$$A = \begin{pmatrix} 2 + 3i & 4 - 5i \\ 3 & 4i \end{pmatrix}$$

$$A^T = \begin{pmatrix} 2 - 3i & 4 + 5i \\ 3 & -4i \end{pmatrix}$$

If  $A^* = A$ , then  $A$  is a **real matrix**.

## Hermitian Conjugate: $A^\dagger$

The **Hermitian conjugate** (or **adjoint**) of an arbitrary matrix  $A$  is obtained by taking the *complex conjugate* of the matrix and then the *transpose* of the complex conjugate matrix.

$$A^\dagger = (A^*)^T = (A^T)^*$$

For example

$$A = \begin{pmatrix} 2 + 3i & 4 - 5i \\ 3 & 4i \end{pmatrix}$$

$$A^\dagger = \begin{pmatrix} 2 - 3i & 4 + 5i \\ 3 & -4i \end{pmatrix}^T = \begin{pmatrix} 2 - 3i & 3 \\ 4 + 5i & -4i \end{pmatrix}$$

It can be shown that

$$(AB)^\dagger = B^\dagger A^\dagger$$

$$(AB \cdots G)^\dagger = G^\dagger \cdots B^\dagger A^\dagger$$

## Inner Product of Vectors

Assume that in a given orthonormal basis the vectors **a** and **b** are represented by the column matrices

$$\mathbf{a} = \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_N \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_N \end{pmatrix}$$

For real vectors

$$\mathbf{a}^T \mathbf{b} = (a_1 \ a_2 \ \cdots \ a_N) \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_N \end{pmatrix} = \sum_{i=1}^N a_i b_i$$

For complex vectors

$$\mathbf{a}^\dagger \mathbf{b} = (a_1^* \ a_2^* \ \cdots \ a_N^*) \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_N \end{pmatrix} = \sum_{i=1}^N a_i^* b_i$$

is the inner product  $\langle \mathbf{a} | \mathbf{b} \rangle$  in that basis.